



The thyroid gland can become overactive (hyperthyroidism) or underactive (hypothyroidism). This is most often due to an autoimmune disease in your body. For more information, visit PreventBlindness.org/thyroid-eye-disease

225 West Wacker Drive
Suite 400
Chicago, Illinois 60606
800.331.2020
PreventBlindness.org


Thyroid Eye Disease

(Also known as Graves' Ophthalmopathy or Graves' Eye Disease)

What is Thyroid Eye Disease?

Thyroid eye disease (TED), also known as Graves' ophthalmopathy or Graves' Eye Disease, is an autoimmune disease in which the immune system causes inflammation and swelling which leads to abnormalities within muscle tissue and fat behind the eye. An overactive thyroid gland (hyperthyroidism) is usually caused by Graves' disease. Up to one-half of people with Graves' disease develop TED. In some people, TED can occur with normal levels of thyroid hormones (euthyroid) or low levels of thyroid hormones (hypothyroidism). TED may occur in patients who already know they have thyroid disease, or it may be the first sign of Graves' disease. While TED often occurs in people living with hyperthyroidism or Graves' disease, it is a distinct disease and treating hyperthyroidism may not resolve the TED symptoms and signs.

The main symptoms of TED include inflammation and an increase in the muscles and fat behind the eye (in the bony eye socket), often causing the eyeballs to push forward. If the eye is pushed far enough forward, the eyelids may not close properly when blinking and sleeping. The clear front part of the eye, called the cornea, may become unprotected, dry, and damaged. Stiffening and inflammation of the muscles of the eye may prevent them from working well, affecting eye position and eye movements and leading to double vision. In severe cases, the inflammation and increase in the tissues, muscles, and fat behind the eye compresses the optic nerve, the nerve that connects the eye to the brain, causing vision loss.



Thyroid Eye Disease

(continued)

Who is at Risk for TED?

TED is most commonly associated with Graves' disease, however it can also occur without a diagnosis of Graves' disease.


Other risk factors for TED include:

- **Age:** Usually affects middle-aged adults but can occur at any age
- **Gender:** Females are affected more than males
- **Family history** of TED
- **Smoking:** Smoking increases the risk of TED by 7–8 times, causes TED to have a longer “active phase”, and it reduces the effectiveness of treatments
- **Radioiodine therapy:** Radioactive iodine has been used to treat hyperthyroidism and Graves' disease. This treatment should be used with caution in people with TED as it may worsen the condition unless steroids are given at the same time
- Low blood levels of **selenium**, a dietary mineral
- Low blood levels of **vitamin D**

What are the Symptoms of TED?

Symptoms of TED are caused by the tissues, fat, and muscles of the eye socket swelling and pushing the eyeball forward. Tissues around the eye also may swell and become thickened. It is not unusual for symptoms to be worse in one eye more than the other. TED changes over time, with symptoms that can appear anytime and range from mild to severe. The appearance of TED can be different for each person.

The symptoms of thyroid eye disease include:



Thyroid Eye Disease

(continued)

In more severe TED, there may also be:

- Trouble moving eyes and closing eyes
- Inability to completely close your eye causing corneal exposure or even an ulcer
- Blurred or loss of vision due to optic nerve compression (colors may appear dull or dim), or corneal damage from an ulcer leading to scarring

Note: If you have difficulty closing your eyelids, you may be at risk to develop a corneal ulcer. The cornea is a clear layer that covers the front of the eye. A corneal ulcer is an open sore on your cornea and it can cause scarring and permanent loss of the vision. A corneal ulcer causes redness of the eye, pain and usually a decrease in vision. You should seek immediate attention from your eye doctor for these problems.

If you have Graves' disease, eye symptoms most often begin within six months of disease diagnosis. Very rarely, eye problems may develop long after the Graves' disease has been treated. In some patients with eye symptoms, hyperthyroidism never develops and, rarely, patients may have hypothyroidism. The severity of the eye symptoms is not related to the severity of the hyperthyroidism.

How is TED Diagnosed?

If you have thyroid disease or suspect that you may have TED, your doctor may recommend that you see an eye doctor (ophthalmologist). TED is managed by an ophthalmologist. Management of more severe forms of the disease may require a team of ophthalmologists, including specialists in eye plastic surgery, eye muscle surgery, and neuro-ophthalmology. They will be able to evaluate your symptoms, recommend a course of treatment, discuss surgical options to improve eye function and appearance, and help you watch for any problems with decreased vision.

To assess your vision and the changes in the tissues around your eyes, there are several tests you can expect your ophthalmologist to perform. The following tests are often conducted:

- **Visual acuity** – to determine how well you can see far away and up close
- **Color vision** – to determine how well you see colors
- **Visual fields** – to assess your central and side vision
- **Eye movements** – to see how much your eye muscles have been affected
- **Eyelid measurements** – to measure the eyelid position on the eye
- **Eye measurements** – to measure the amount of eye bulging (protrusion)

Thyroid Eye Disease

(continued)

- **Eye pressure readings** – to determine the pressure inside the eye
- **Optic nerve visualization** – to look into the back of your eye to see if the optic nerve is affected
- **Magnetic resonance imaging (MRI) scan or computed tomography (CT) scan** – to see how much enlargement there is in the tissue and muscles behind the eye

How is TED Treated/Managed?

TED in its severe form can last between one and three years. That means if it is left untreated, the inflammation may gradually decrease by itself but may cause damage to vision through the course of the disease. Inflammation also can flare up later. Sometimes, the changes caused by the enlargement of the tissues (such as bulging eyes or double vision) may not go away. The goal of treatment is to limit severe inflammation and swelling and to protect the front of the eye and prevent vision loss.

TED is managed by an ophthalmologist. Any underlying thyroid problems will be managed by your primary care doctor (PCP) or by a specialist in the hormone systems of the body (an endocrinologist).

If a thyroid issue is suspected, evaluation and treatment are critical. The first priority is to restore your normal thyroid function. In addition, eye conditions should be examined and treated at the same time as your thyroid gland treatment. Eye problems may continue to progress even after your thyroid function returns to normal.

What Can I Do to Help My Vision?

If you have TED, your ophthalmologist may recommend one or more of the following treatments to help soothe your eyes and improve your vision:

- **Cool compresses:** Apply cool compresses to your eyes. The extra moisture and cooling effect may provide relief.
- **Sunglasses:** When you have TED, your eyes are more sensitive to sunlight and UV rays. Wearing sunglasses helps protect your eyes from both sun and wind.
- **Lubricating eye drops:** Use lubricating eye drops, like artificial tears. It may help relieve dryness and scratchiness. Make sure to use eye drops that do not contain redness removers. Lubricating gels can be used before bed to prevent the cornea (the front of the eye) from drying out because your eyelids may not close completely when sleeping.
- **Taping:** Talk with your doctor about taping your top and bottom eyelids together to help protect your front of your eye (cornea) from drying when your eyelids do not close completely during sleep.

Thyroid Eye Disease

(continued)

- **Elevate your head when lying down:** Keeping your head higher than the rest of your body may reduce swelling and may help relieve pressure on your eyes.
- **Quit smoking:** Smoking (and secondhand exposure to smoke) is an important risk factor for TED. If you smoke, quit, and avoid secondhand smoke.
- **Steroids:** Swelling in your eyes may be improved by treatment with steroids (such as hydrocortisone or prednisone). Your ophthalmologist may recommend either intravenous or oral medication. Note, make sure you discuss the risks of use of steroids with your doctor before use.
- **Pharmaceutical treatments:** There are intravenous infusion (putting drugs into a vein) treatments that are FDA-approved medications for the treatment of TED. Researchers are studying new ways to treat TED and several other treatments are in clinical trials. If you're interested in learning more or seeing if a trial might be right for you, talk to your doctor or visit clinicaltrials.org. You can learn more about clinical trials at: PreventBlindness.org/clinical-trials-for-eye-diseases-and-vision
- **Selenium supplements:** More research is needed around the effectiveness of selenium supplements. Talk to your doctor before starting any supplements.
- **Prisms:** TED can cause scar tissue to develop in your eye muscles. This can lead them to become tight and pull your eyes out of alignment, causing double vision. If double vision occurs, glasses containing prisms may be prescribed by your doctor. However, prisms do not work for all people with double vision and your doctor may recommend patching one eye for temporary relief or eye muscle surgery as a more effective option when changes have stabilized.
- **Eyelid surgery:** When you have TED, the eyelids are usually more widely open with a “startled” look because the muscles in the eyelids may tighten and pull the upper lid up and the lower lid down. You may have difficulty closing your eyelids, leaving the front of the eye (cornea) more exposed, which causes tearing, irritation and susceptibility to developing a corneal ulcer. Eyelid surgery may help reduce exposure of the cornea.
- **Eye muscle surgery:** Eye muscle surgery may help correct your double vision by moving the affected muscle(s) farther back from its original position on the eyeball. This surgery will help to correct your double vision when reading and looking straight ahead. Sometimes, you may need more than one surgery to get effective results.

Thyroid Eye Disease

(continued)



- **Orbital decompression surgery:** TED can cause swollen tissues around the eye that compresses the optic nerve. The optic nerve provides the connection between your eye and the brain. When the nerve is compressed, color vision becomes abnormal, lights may seem dimmer than usual, and the sharpness of the vision decreases. Orbital decompression surgery can be done to improve your vision. The surgery makes the eye socket bigger or removes some of the excess tissue. When the nerve is compressed, the goal of surgery is to get the eye and the inflamed tissue more space and decrease pressure on the optic nerve. Even when the optic nerve function is not compromised, orbital decompression may be used to restore comfort and appearance by reducing the bulging of the eyes. If orbital decompression surgery is recommended, it is usually performed prior to eye muscle surgery and/or eyelid surgery, if needed.

Your ophthalmologist will require some time for your TED to stabilize before recommending surgery. Typically, the active or inflammatory stage of TED lasts one to three years. During this time, your ophthalmologist will avoid surgically treating your symptoms unless your vision is threatened. In cases such as a corneal ulcer or optic nerve compression urgent surgery may be recommended.

How to support your coverage options with TED?

Reach out to your insurance company directly: This is the most important step. Describe your circumstances, including your TED diagnosis, and ask if they offer coverage for health services.

Understand your insurance processes/requirements: Every insurance carrier is different. Request information directly from them about the process for securing coverage, including any requirements for a Letter of Medical Necessity (LMN) from a physician.

Talk to your human resources department: Look into benefits your employer provides. Some employers provide wellness initiatives or perks that could encompass coverage or reimbursement for health needs like emergency funds to cover for expensive treatments.

How Does TED Affect Mental Health?

TED can cause eye pain, double vision, loss of vision, and changes to your appearance which may impact your quality of life. The impacts of TED may lead you to experience depression, anxiety, loss of independence, and reduced self-confidence. You may have a decreased desire to socialize with others, have trouble with productivity at school or work, and stop doing activities you once enjoyed.

Thyroid Eye Disease

(continued)

To help support your mental health, consider the following:

- **Talk to your doctor:** Talk to your doctor about any changes to your emotional well-being. Ask questions on how TED will affect your daily life and work.
- **Seek support:** Seek out support groups of others who are going through TED. The TED Community Organization (tedcommunity.org/) and The Graves' Disease and Thyroid Foundation ([GDATE.org/community/support-groups](https://gdatf.org/community/support-groups)) offers support groups for people with Graves' disease, TED, and other thyroid conditions. These groups can help you learn about your condition, share your experience, and find support.
- **Connect with others:** Stay connected with friends and family to keep you from feeling isolated. Consider involving your closest family and friends in your treatment, let them know how they can support you during this process.
- **Take time to enjoy life:** Take part in activities or hobbies that make you happy.
- **Exercise:** Exercise can help symptoms of depression or anxiety and make you feel better. Talk to your health care professional to determine what exercise routine may work best for you, especially if you have changes to your vision due to TED.

To learn more, check out the following resources:

National Alliance on Mental Illness (NAMI):

Warning signs and symptoms [NAMI.org/About-Mental-Illness/Warning-Signs-and-Symptoms](https://nami.org/About-Mental-Illness/Warning-Signs-and-Symptoms)

American Psychological Association: Depression [APA.org/topics/depression](https://apa.org/topics/depression)

Centers for Disease Control and Prevention (CDC): The Mental Health of People with Disabilities

cdc.gov/disability-and-health/articles-documents/mental-health-of-people-with-disabilities.html

Vision Loss and Mental Health cdc.gov/vision-health/about-eye-disorders/vision-loss-mental-health.html

For more information, visit PreventBlindness.org/thyroid-eye-disease

TED community organization tedcommunity.org

Focus On Eye Health Expert Series: Thyroid Eye Disease (TED)

youtu.be/-g2UPKrQWCU?feature=shared

Made possible by funding from:

